## PREPARATION FOR COLONOSCOPY MORNING PROCEDURE

In order to ensure the best possible outcome with respect to your colonoscopy, it is essential that the large bowel is completely clear of waste material. If the bowel is not clear, the procedure may have to be repeated. It will be necessary to modify your diet for a couple of days prior to colonoscopy and to take a special mixture prior to your procedure.

Iron tablets should be stopped 7 days prior to the procedure. If you are taking Plavix, Iscover, Effient, Brilinta or Warfarin or if you are a Diabetic, please ensure you advise our rooms at least 7 days prior to the procedure day.

If you feel you have a health problem which will affect your ability to follow the directions, please contact my rooms. The regimen for your preparation is as follows:

DAY	AVOID	ALLOWED	
Day 1	High fibre/high residue foods:		
2 days before your procedure	<ul> <li>Most breakfast cereals</li> </ul>	<ul> <li>White bread (no added fibre)</li> </ul>	
	<ul> <li>Non-white bread (rye,</li> </ul>	<ul> <li>White flour products</li> </ul>	
	wholemeal, multigrain)	<ul> <li>White rice</li> </ul>	
	<ul> <li>Grains &amp; legumes</li> </ul>	<ul> <li>Rice Bubbles &amp; Cornflakes</li> </ul>	
	<ul> <li>Nuts &amp; Seeds</li> </ul>	<ul> <li>Dairy products – milk, cheese – up to 2</li> </ul>	
	<ul><li>Salads</li></ul>	cups	
	<ul> <li>Uncooked fruit and</li> </ul>	<ul> <li>Fish, eggs and chicken</li> </ul>	
	vegetables	<ul> <li>Peeled and mashed potato or pumpkin</li> </ul>	
	<ul> <li>Green vegetables</li> </ul>	<ul> <li>Peeled stewed fruit</li> </ul>	
	<ul> <li>Red meat</li> </ul>		
Day 2		A light breakfast such as tea and toast, then:	
1 day before your procedure		<ul> <li>2 glasses of <u>strained</u> fruit juice for the</li> </ul>	
	After breakfast :	eakfast : day	
	<ul> <li>NO solid food</li> </ul>	<ul> <li>Clear fruit juice such as apple (not</li> </ul>	
	<ul> <li>NO Red, green or</li> </ul>	cloudy)	
	purple jelly or cordial	<ul> <li>Tea or coffee with a <u>small amount of</u></li> </ul>	
	<ul> <li>NO Dark grape juice or</li> </ul>	<u>milk</u>	
	blackcurrant juice	<ul> <li>Orange or yellow jelly and cordial</li> </ul>	
		<ul> <li>Lemonade or soft drinks (not red or</li> </ul>	
		green)	
		<ul> <li>Clear soup or fluids</li> </ul>	
		<ul> <li>100 g tub of plain yoghurt</li> </ul>	
		<ul> <li>1 small scoop of vanilla icecream</li> </ul>	
		UP UNTIL 4.00 PM	

VOLIR PREPARATION IS:	2 sachets of Picolax and 2	sachets of Coloni VTFI V	and the instructions are:

On DAY 2, Between 4:00 & 6.00pm Take 1 sachet of Picolax, as per packet instructions + 1 extra glass of water.

- **1 Hour later** Dissolve 1 sachet of ColonLYTELY in 1 litre of water and drink over the next hour i.e. 1 x 250 ml glass every 15 minutes. Refrigerate the mixture if you prefer.
- 1 Hour after finishing the colonLYTELY take the second sachet of Picolax as per the packet instructions.

  Consume another 2 glasses of water before bed

Sorry, you will need to take more preparation early in the morning!	Take 1 litre of colonLYTELY over 1 hour, ie. 250mls every 15 minutes, commencing 3 hours before your appointment time.	ABSOLUTELY NIL BY MOUTH
		FROM
Taking preparation up to 2 hours	As your time of arrival at the procedure centre is	FROIVI
before your appointment time has	Commence the colonLYTELY at	·
been shown to result in a much		
cleaner bowel and this is very		
important.		

## **Helpful Hints**

- 1. Suck a peppermint between each glass of ColonLYTELY
- 2. Apply a protective cream e.g. lanolin to your bottom to reduce scalding.
- 3. Wear loose, casual clothing when you come along for the colonoscopy.

If you have any queries, please call our rooms and the staff will be happy to assist you Ph: 07 5591 4455

> Mon – Thurs 08:30 am – 4:30 pm Friday 08:30 am – 4:00 pm